

# Meningococcal Disease

## What is meningococcal disease?

Meningococcal disease is a rare but serious illness caused by bacteria (germs) called *Neisseria meningitidis*. The different strains ('serogroups') of these bacteria are named by letters of the alphabet such as A, B, C, W and Y.

The most common strain has changed over time. Meningococcal C vaccine became part of the National Immunisation Program in 2003, and disease due to strain C is now rare.

Throughout Australia most meningococcal disease is now caused by strains W and B. Strain W increased significantly throughout Australia from 2014 onwards.

## What are the symptoms?

### ***Meningitis (inflammation of the membranes covering the brain and spinal cord)***

- Fever
- Headache
- Neck pain or stiffness
- Pain from lights
- Drowsiness or confusion.

### ***Septicaemia (blood infection)***

- Fever
- Muscle aches
- Vomiting
- Rash (a red rash which does not fade when pressed and can become dark purple).

## Other symptoms

- Joint pain and swelling
- Lung infection with fever, cough, chest pain and difficulty breathing
- Severe throat infection with pain, difficulty swallowing, and drooling.

Children and adolescents can have leg pain, cold hands and feet, and blotchy skin.

Babies and infants often don't have these symptoms but can be unsettled or drowsy, floppy and not feeding.

It can take from one to seven days from being exposed to the bacteria to becoming sick. Symptoms can become life threatening within hours.

## How is it spread?

Meningococcal bacteria are spread in secretions of the throat and nose. This is more likely to happen among people with close, regular and prolonged contact.

About 1 in 10 people can have meningococcal bacteria in their throat or nose. These very rarely cause illness.

Regular exposure to tobacco smoke can increase the risk of catching the bacteria and becoming ill.

The bacteria don't survive outside the body and don't spread easily from shared drinks, food or cigarettes.

## How is it diagnosed?

Meningococcal disease is diagnosed in blood samples, or fluid samples from around the spinal cord.

## How is it treated?

People with meningococcal disease need urgent treatment in hospital. Treatment usually starts before test results are known. Treatments include:

- Antibiotics
- Medications for fever and pain
- Fluids

A person is no longer contagious after 24 hours of antibiotics.

## How is it prevented?

Immunisation is the best way to prevent meningococcal disease.

### Meningococcal ACWY vaccine

From 1 July 2018 a quadrivalent meningococcal vaccine for strains A, C, W and Y replaced the meningococcal C vaccine as part of the National Immunisation Program for all children aged 12 months.

All Tasmanian teenagers who were born between 1 August 1997 and 30 April 2003 are eligible to receive a free meningococcal W vaccine. The vaccine is available through general practitioners and some council clinics.

ACWY vaccine is also available on private prescription for those who wish to protect themselves or their family from these strains of meningococcal disease.

**Meningococcal B vaccine** is available on private prescription.

- Recommended for infants and young children, adolescents, young adults living close together (such as students in residential accommodation), some medical conditions and occupations.

- Can also be offered to those who wish to protect themselves or their family from this strain of meningococcal disease.

## What should I do if I have had contact with someone who has meningococcal disease?

Antibiotics are given to people (contacts) who live in the same house, and those who have spent a long time or been intimately close, with someone with the disease. This is because one of these people may have the bacteria in their throat or nose. Antibiotics 'clear' the bacteria out of their throat and nose so they can't pass it to other people. Contacts may also be offered a meningococcal vaccine.

It is rare for people to develop the illness after being in close contact with someone who has meningococcal disease, but you should still watch out for symptoms of meningococcal disease.

Contacts of someone who has meningococcal disease can go to work, school and childcare.

## What should I do if I have meningococcal disease?

If you think you or someone close to you has meningococcal disease seek help immediately from your doctor, hospital emergency department or ambulance.

### More information

[www.health.gov.au/internet/main/publishing.nsf/Content/ohp-meningococcal-WV.htm](http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-meningococcal-WV.htm)

<http://www.ncirs.edu.au/provider-resources/ncirs-fact-sheets/>

Public Health Hotline – Tasmania on **1800 671 738** to speak to a clinical nurse consultant

*July 2018*